

A RESOLVED DEBRIS DISK AROUND THE G2 V STAR HD 107146

D. R. ARDILA,¹ D. A. GOLIMOWSKI,¹ J. E. KRIST,² M. CLAMPIN,³ J. P. WILLIAMS,⁴ J. P. BLAKESLEE,¹
H. C. FORD,¹ G. F. HARTIG,² AND G. D. ILLINGWORTH⁵

Received 2004 August 25; accepted 2004 November 15; published 2004 November 22

ABSTRACT

We present resolved scattered-light images of the debris disk around HD 107146, a G2 star 28.5 pc from the Sun. This is the first debris disk to be resolved in scattered light around a solar-type star. We observed it with the *Hubble Space Telescope*/Advanced Camera for Surveys coronagraph, using a 1"8 occulting spot and the F606W (broad *V*) and F814W (broad *I*) filters. Within 2" from the star, the image is dominated by point-spread function subtraction residuals. Outside this limit, the disk looks featureless, except for a northeast-southwest brightness asymmetry that we attribute to forward scattering. The disk has scattered-light fractional luminosities of $(L_{\text{sca}}/L_*)_{\text{F606W}} = (6.8 \pm 0.8) \times 10^{-5}$ and $(L_{\text{sca}}/L_*)_{\text{F814W}} = (10 \pm 1) \times 10^{-5}$, and it is detected up to 6"5 away from the star. To map the surface density of the disk, we deproject it by $25^\circ \pm 5^\circ$, divide by the dust-scattering phase ($g_{\text{F606W}} = 0.3 \pm 0.1$, $g_{\text{F814W}} = 0.2 \pm 0.1$), and correct for the geometric dilution of starlight. Within the errors, the surface density has the same shape in each bandpass, and it appears to be a broad (85 AU) ring with most of the opacity concentrated at 130 AU. The ratio of the relative luminosity in F814W to that in F606W has the constant value of 1.3 ± 0.3 , with the error dominated by uncertainties in the value of g in each filter. An examination of far-infrared and submillimeter measurements suggests the presence of small grains. The colors and the derived values of g are consistent with the presence of dust particles smaller than the radiation pressure limit. Possible scenarios that may explain the shape of the surface density profile are the creation of a small planet or the scattering and circularization of a large one.

Subject headings: planetary systems: formation — planetary systems: protoplanetary disks — stars: imaging — stars: individual (HD 107146)

Online material: FITS files

1. INTRODUCTION

The presence of dusty disks around main-sequence stars serves as a marker for the existence of planetesimals. Without collisions among planetesimals, or their evaporation, the dust would not be replenished, and it would have disappeared from the system long ago. Thus, debris disks indicate that the planet formation process is occurring or has occurred. In particular, the study of disks around low-mass stars illuminates the planet formation process in the relatively low radiation environments analogous to the solar system. Resolved images of these systems help to constrain their physical and geometrical properties. Scattered-light images sample the whole disk regardless of its temperature, and this, coupled with the fact that optical and near-infrared detectors have higher angular resolution than far-infrared and submillimeter ones, allows us to have a rich understanding of the systems.

Here we present coronagraphic scattered-light images of the disk around the G2 V star HD 107146. This is the first debris disk resolved in scattered light around a solar-like star. The observations presented here address the issue of the diversity of planet formation histories within stars of the same spectral type, like HD 107146 and our Sun. To date, only one other debris disk around a non-A star (the M0 star AU Mic; see Krist et al. 2004, Liu 2004, and Kalas et al. 2004) has been resolved in scattered light.

The disk around HD 107146 was marginally resolved by Williams et al. (2004) at 450 μm . Submillimeter and mid-infrared measurements indicate that its fractional excess luminosity is $f_d = L_d/L_* = 1.2 \times 10^{-3}$, due to a dust mass of 0.1–0.4 M_\oplus , comparable to that of the β Pictoris disk. An inner hole (>31 AU) is suggested by the lack of *IRAS* 25 μm detection. No gas measurements have been reported. The *Hipparcos* distance of the host star is 28.5 pc, its luminosity is 1.1 L_\odot , and age estimates range from 30 to 250 Myr (Williams et al. 2004). The star is a candidate periodic *V*-band photometric variable ($P = 7$ days; Koen & Eyer 2002).

2. OBSERVATIONS

HD 107146 and the point-spread function (PSF) reference star HD 120066 were observed with the *Hubble Space Telescope* (*HST*) Advanced Camera for Survey (ACS) high-resolution camera (HRC) on UT 2004 June 5 and July 20. The observations were conducted as part of the guaranteed observing time awarded to the ACS Investigation Definition Team (proposal IDs 9987 and 10330). The images were taken with the F606W (broad *V*) and F814W (broad *I*) filters. For each band and each target, a short direct exposure was followed by a coronagraphic exposure using the 1"8 mask. The coronagraphic exposures for the target were 2330 s long in F606W and 2150 s long in F814W. For HD 120066, they were 1990 s in F606W and 2250 s in F814W.

Here we present a summary of the reduction procedure: a more detailed description, applied to different targets, can be found in Krist et al. (2004) and Clampin et al. (2003). The results of the measurements are in Table 1. To measure the magnitudes of the target and the PSF reference star, we integrated the direct image flux within a circular aperture greater

¹ Department of Physics and Astronomy, Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218.

² Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218.

³ NASA Goddard Space Flight Center, Code 681, Greenbelt, MD 20771.

⁴ Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, HI 96822.

⁵ UCO/Lick Observatory, University of California, Santa Cruz, CA 95064.

TABLE 1
MEASURED QUANTITIES

Band	Stellar Brightness	L_{scat}/L_* ^a	Disk Magnitude ^b	g^c
F606W	5.9 ± 0.3 Jy, $V = 7.05 \pm 0.05$	$(6.8 \pm 0.8) \times 10^{-5}$	$V = 17.5 \pm 0.1$	0.3 ± 1
F814W	6.9 ± 0.3 Jy, $I = 6.35 \pm 0.05$	$(10 \pm 1) \times 10^{-5}$	$I = 16.4 \pm 0.1$	0.2 ± 1

^a Measured in a circular annulus between $3''.5$ and $6''.0$ (the brightest part of the disk).

^b Integrated magnitude in a circular annulus between $3''.5$ and $6''.0$.

^c Scattering asymmetry parameter in a Henyey-Greenstein phase function.

than $6''$ in radius, which includes the saturated stellar core.⁶ The transformation between counts per second and magnitudes was obtained using the STSDAS synthetic photometry package SYNPHOT, which simulates most *HST* observing configurations. The coronagraphic image of HD 120066 was normalized to, aligned with, and subtracted from the image of HD 107146. Alignment of the images is accurate to within ± 0.2 pixels ($\pm 0''.005$). The resulting subtracted image was smoothed using a 3×3 median filter and corrected for geometric distortion. Figure 1 shows the result. To increase the signal-to-noise ratio of the displayed images, we have performed an additional 5×5 median smoothing and rebinned by a factor of 2.

Subtraction errors, caused by mismatches in the colors of the stars or PSF time variability, dominate the emission within $\sim 2''$ from the star and contribute light at large distances. With only one reference star, we cannot quantify very precisely the magnitudes of these two error sources. The ACS Instrument

⁶ Instrument Science Reports, R. L. Gilliland, 2004 January 5, <http://www.stsci.edu/hst/acs/documents/isrs>.




FIG. 1.—Disk around the G2 V star HD 107146. *Top left*: F606W filter (broad V); *top right*: F814W filter (broad I). To obtain these images, a reference PSF (HD 120066) was normalized to, aligned with, and subtracted from the coronagraphic images of HD 107146. The $3''.0$ spot in the upper right-hand corner has been masked out. *Bottom*: Maps of the scattering optical depth τ_{ω} (the product of the total optical depth times the albedo), obtained by de-projecting the disk (assuming an inclination of 25° from face-on), multiplying by the distance squared from the stellar position, correcting for forward scattering, and dividing by the stellar flux. *Bottom left*: F606W filter; *bottom right*: F814W filter. Note that the bottom row images have the same color scale but that the top row ones do not. FITS versions of these images are available as a gzipped tar file in the electronic edition of the Journal.

Handbook⁷ indicates that a mismatch of three spectral classes would produce subtraction residuals for a star of this brightness of the order of $0.2 \mu\text{Jy arcsec}^{-2}$ (or $25.73 \text{ mag arcsec}^{-2}$ in the V band), $5''$ away from the target. The actual errors in these observations are likely to be smaller because the $V-I$ colors of the two stars are the same within errors: for HD 120066, we measure $V = 6.32 \pm 0.05$, $V - I = 0.68 \pm 0.07$. The instrument handbook also indicates that typical time-dependent PSF variations within an orbit (due to variations in focus) will result in errors of the same order. However, Figure 2 suggests that this may underestimate the error in our observations, as subtraction residuals $6''.5$ from the star occur at the level of $\sim 1 \mu\text{Jy arcsec}^{-2}$.

To quantify the systematic error in the normalization constant between the target and the PSF reference star, we compare the value of this constant obtained by using four different methods: taking the ratio of the stellar flux in each band, taking the ratio of the flux of each star away from the saturated columns, taking the ratio of the number of saturated pixels per unit time in the direct images for each band, and adjusting the value of the constant to produce the cleanest visual PSF subtraction. For each filter, the four methods yield the same normalization constant within 1%, which corresponds to photometric errors of 0.1 and 0.2 mag arcsec⁻² in the brightest and faintest regions of the disk, respectively. Given that this is an estimate of a systematic error, in what follows we propagate it linearly to

⁷ ACS Instrument Handbook, <http://www.stsci.edu/hst/acs/documents/handbooks/cycle13/cover.html>.




FIG. 2.—Surface brightness as a function of distance, obtained by taking the median of the flux in rectangular boxes $2''$ wide (centered in the stellar position) by $0''.125$ long. The measurements are taken along a path with a position angle of 58° (the disk's minor axis). Positive displacements are toward the southwest. The thick-black (red-dashed) line is the trace in the F606W (F814W) band. The dotted lines are positioned at $2''$ from the star. This profile suggests that subtraction residuals dominate beyond $6''.5$, at the level of $\sim 1 \mu\text{Jy arcsec}^{-2}$; $10 \mu\text{Jy arcsec}^{-2}$; $10 \mu\text{Jy arcsec}^{-2}$ correspond to $V = 21.48$ or $I = 20.95$.

estimate uncertainties in calculated quantities. For the quantities calculated below, this error dominates over all others, including the $\sim 5\%$ random photometric error.

3. RESULTS

The circumstellar disk is clearly seen in the subtracted images. Within the limitations of the observations, it appears elongated along the southeast-northwest direction and featureless, except for the fact that the southwest side is brighter than the northeast side. The excess color of the disk with respect to the star is $\Delta(V - I) = 0.4 \pm 0.3$ (Table 1). This would be the intrinsic color of the disk if the scattering phase function were independent of wavelength. We also detect an object 6''8 southwest of the star. The time baseline between the observations taken in the two filters is too short to detect any difference in the relative positions between the star and the object. Smooth elliptical fits subtracted from the object reveal residual spiral structure. Therefore, we believe this to be a faint background spiral galaxy ($V = 19.4$, $V - I = 1.2$). The galaxy is at a different position angle than the offset with respect to the optical position of the star of the SCUBA 450 μm or the map's extension (Williams et al. 2004). This suggests that the submillimeter measurements are not contaminated by the presence of the galaxy.

The slightly red color of the disk with respect to the star suggests the presence of small grains, which leads us to interpret the northeast-southwest brightness asymmetry as being due to preferential forward scattering in an inclined disk. This interpretation is consistent with the slightly elongated shape of the observed disk. By fitting elliptical isophotes to the disk image, we conclude that the disk minor axis has a position angle of $58^\circ \pm 5^\circ$. The bright southwest region is symmetric with respect to this axis. The measurements are consistent with the picture of a circular disk inclined $25^\circ \pm 5^\circ$ from the plane of the sky.

Assuming that the disk is optically thin (as implied by its low f_d -value), we can map the surface density (Clampin et al. 2003). We deproject the disk, assuming that it is intrinsically circular, and multiply the resulting image by r^2 , where r is the distance to the star, to correct for the geometric dilution of starlight. Finally, we divide the deprojected image by a Henyey-Greenstein phase function, adjusting g , the scattering asymmetry parameter, until the front- and back-scattering regions have the same brightness (Table 1). After dividing by the stellar luminosity, the result (Fig. 1, *bottom panels*) is a map of the scattering optical depth, which we write as $\tau\omega$, the optical depth times the albedo in the band. The scattering optical depth is proportional to the surface density (Weinberger et al. 1999; Clampin et al. 2003). To eliminate the galaxy, we fitted its isophotes to a series of ellipses of varying centers, ellipticities, and position angles and generated a model to be subtracted from the image.

Within the level of the errors, the disks shown in the bottom panels of Figure 1 are azimuthally featureless, and they have the same shape. The observed morphology is a broad ring, with maximum opacity at 130 AU and a FWHM of 85 AU. By taking medians of annular sections of the deprojected disks, we map the scattering optical depth as a function of distance (Fig. 3, *top panel*). The shape of $\tau\omega$ can be parameterized, as r^p , by two power laws: $p = 1.6 \pm 0.5$ (from 80 to 130 AU) and $p = -2.8 \pm 0.3$ (from 130 to 185 AU). Given the large subtraction residuals within ~ 60 AU from the star, our observations are not inconsistent with the presence of dust within this radius or with a constant optical depth within 80 AU. However, we clearly detect a decrease in the dust opacity within 130 AU: the normalization error would have to be larger than




FIG. 3.—*Top*: Median $\tau\omega$ as a function of distance from the star, obtained in a series of annuli 0".14 wide, measured from the bottom panels of Fig. 1. The solid line is the F606W observation ($g_{\text{F606W}} = 0.3 \pm 0.1$). The red-dashed line is the F814W observation ($g_{\text{F814W}} = 0.2 \pm 0.1$). The error bars indicate the range of values obtained if the standard PSF is oversubtracted or under-subtracted by 1%. *Bottom*: Ratio of $(\tau\omega)_{\text{F814W}}$ to $(\tau\omega)_{\text{F606W}}$, as a function of distance from the star. As above, the error bars show the dependence on errors in the normalization. The dashed lines indicate the dependence of the ratio on the value of g : the top dotted line is obtained assuming $g_{\text{F606W}} = 0.2$ and $g_{\text{F814W}} = 0.3$, and the bottom one is obtained assuming $g_{\text{F606W}} = 0.4$ and $g_{\text{F814W}} = 0.1$. These choices produce extremal ratios.

calculated ($>5\%$) for the observations to be consistent with constant opacity within this limit. The difference in the shape of $\tau\omega$ between the two bands is not significant, given the subtraction residuals in the deprojected F814W image.

For the adopted values of g , the scattering optical depth is similar in both bands. To measure the color of the deprojected disk, we take the ratio of the two deprojected images and obtain medians of annular sections of the result (Fig. 3, *bottom panel*). The color can be reliably determined only between 100 and 180 AU. Between these limits the disk is uniform in color, with the mean of the ratio in $\tau\omega$ of the two filters being 1.3 ± 0.3 . The uncertainty is determined from the uncertainty in the g -values. The uncertainty in the mean is ± 0.06 .

4. ANALYSIS AND DISCUSSION

The outer radius of the disk resolved in scattered light is similar to the submillimeter one. To fit the thermal emission, Williams et al. (2004) used a single-temperature modified blackbody function, in which the emission is proportional to $Q_\lambda B_\lambda$, with $Q_\lambda = 1 - \exp[-(\lambda_0/\lambda)^\beta]$ and $\lambda_0 = 100 \mu\text{m}$ (the characteristic grain size). They concluded that $T = 51 \pm 4$ K and $\beta = 0.69 \pm 0.15$.

A detailed model of the thermal infrared emission of the system is beyond the scope of this Letter. Here we note that if we constrain the thermal emission to originate at 130 AU (the radius of maximum brightness in scattered light) and use the values of λ_0 , β , and T from Williams et al. (2004), the disk will not be in thermal equilibrium with the stellar radiation (Backman & Paresce 1993, eq. [1]). In the context of this model, the only way to preserve thermal equilibrium is by reducing λ_0 , the characteristic particle size. By keeping $\beta = 0.7$ constant, we fit the data with $T \sim 45 \pm 5$ K and $\lambda_0 = 2 \pm 1 \mu\text{m}$. With $\beta = 1$, we obtain $T \sim 40 \pm 5$ K and $\lambda_0 = 15 \pm 3 \mu\text{m}$. The precise values of β and λ_0 are poorly constrained, as shown by the fact that both sets of parameters fit the thermal data and satisfy thermal equilibrium. For comparison, Dent et al. (2000) found that for

β Pic, ϵ Eri, Vega, and Fomalhaut, $\lambda_0 = 10\text{--}100 \mu\text{m}$ and $\beta = 0.8\text{--}1.1$. The $25 \mu\text{m}$ *IRAS* nondetection (Williams et al. 2004) and the results of the fit (with $\beta = 0.7$) imply that the dust surface density at 130 AU is ≥ 5 times more than at 60 AU, similar to the conclusion reached in scattered light.

Even though it is poorly constrained, the value of λ_0 suggests the presence of small grains in the disk. The scattering asymmetry parameter, $g = 0.2\text{--}0.3$, is also consistent with the presence of small grains. Similar values of g are obtained in other debris disks: e.g., $g = 0.15\text{--}0.25$ for HD 141569A⁸ (Clampin et al. 2003) and $g = 0.4$ for AU Mic (Krist et al. 2004). For the standard “astronomical silicate” (Draine & Lee 1984; Laor & Draine 1993), such values indicate the presence of submicron grains, although the actual predicted size is not very sensitive to the value of g (Weingartner & Draine 2001). The color is also sensitive to grain size. Assuming compact astronomical silicate particles with size distribution $s^{-3.5}$, where s is the particle radius, one can obtain a color ratio between the two bands as large as ~ 1.2 if the lower limit of the size distribution is $\sim 0.3 \mu\text{m}$. With this dust model, a gray disk is observed when the dust particles are $\geq 1 \mu\text{m}$. The ratio of the force due to radiation pressure to the gravitational force is $\beta_{\text{RP}} = 0.5/s$, with s in microns (assuming a unity radiation pressure coefficient and a dust density of $\rho_d = 1.25 \text{ g cm}^{-3}$; see Takeuchi & Artymowicz 2001). Grains with $\beta_{\text{RP}} > 0.5$ or $s < 1 \mu\text{m}$ will be expelled from the system in dynamical timescales. In other words, the mean color is consistent with the presence of grains smaller than the radiation pressure limiting size. A more detailed analysis (with a more realistic dust model) is necessary to confirm this result, although a similar situation has been found for the debris disk around HD 141569 (Ardila et al. 2005).

The appearance of a symmetric ring is one of the features of the models by Takeuchi & Artymowicz (2001). Another is dust segregation: their models predict that the smallest grains present in the system ($\beta_{\text{RP}} \sim 0.5$) should reside at the largest radii and that larger grains should remain closer to the star. The actual parameters of the segregation depend on the behavior of the gas density at the disk edge. Even in the absence of gas, smaller grains should, on average, reside farther out than larger grains. Over the wavelength span of the two bands, we detect no significant difference in the surface density profile or any systematic color change as a function of distance. Comparisons among observations performed over a wider range of wavelengths (e.g., NICMOS vs. ACS/HRC) are crucial to detect any size separation.

Kenyon & Bromley (2004) show that increased collisions among planetesimals due to the formation of planets with radii larger than 1000 km can generate dust rings. In their models,

⁸ Because of a typo, the value of g for the HD 141569A disk is quoted as $g = 0.25\text{--}0.35$ in Clampin et al. (2003).

the outer edge of the ring marks the position at which planets are starting to form. From their calculations, a planet at 170 AU could grow to the appropriate size if the mass in solids of the disk is between ~ 10 and ~ 70 times larger than that of the minimum-mass solar nebula. The range is due to the uncertainty in the age of HD 107146. The dust ring maximum fractional luminosity would be $f_d \sim 10^{-2}$, larger than the observations. On the other hand, these kind of models produce very axisymmetric structures, similar to those seen in Figure 1.

An alternative to local planet formation is the migration of a planet formed at a smaller radius. Weidenschilling & Marzari (1996) show that gravitational interactions among multiple large bodies can scatter one of them into an eccentric large orbit in dynamical timescales. Dynamical friction will induce large eccentricities in smaller bodies, increasing their collision rate and generating broad rings (Kenyon et al. 1999). The timescale for this process will depend on the relative masses of the planet and the planetesimals. This scenario would imply that at least two giant planets are present in the HD 107146 system and that the presence of the hole reflects the relative paucity of an underlying population of planetesimals, expelled by the (giant) planets. A 50 Myr old, $10M_{\text{Jup}}$ planet would have $m_1 \sim 23 \text{ mag}$ (Burrows et al. 1997). If the PSF is spread over 4 HRC pixels, each would have a brightness of $16.5 \text{ mag arcsec}^{-2}$. In the F814W-subtracted image of HD 107146, the mean surface brightness within $1''.8$ is $15.9 \pm 2 \text{ mag arcsec}^{-2}$, which implies that the planet would not be detected photometrically.

Does this disk represent an earlier stage in the evolution of our solar system? The observed dust disk is larger and proportionally much wider than the solar system Kuiper Belt (KB), and it has ~ 4 orders of magnitude more dust (Greaves et al. 2004). In order to look like the KB, the disk would have to shrink by a factor of ~ 3 and become narrower by a factor of ~ 8 . This evolutionary path seems contrary to current ideas about the solar system. Gomes (2003) argues that the KB formed closer inward and was pushed out by interactions with Neptune. Additionally, Levison & Morbidelli (2003) argue that the sharp exterior edge of the KB is determined by a 2 : 1 resonance with Neptune. Defining the edge of the HD 107146 disk at 185 AU implies a planet at 116 AU. There is no dynamical or photometric signature of this object in the scattered-light image. We believe therefore that HD 107146 is unlikely to evolve into a system like our own.

We wish to thank the team from Williams et al. (2004) for providing us with the name of their target before publication. ACS was developed under NASA contract NAS 5-32865, and this research has been supported by NASA grant NAG5-7697. We are grateful for an equipment grant from Sun Microsystems, Inc. The Space Telescope Science Institute is operated by AURA, Inc., under NASA contract NAS5-26555.

REFERENCES

- Ardila, D. R., et al. 2005, *ApJ*, submitted
 Backman, D. E., & Paresce, F. 1993, *Protostars and Planets III*, ed. E. H. Levy & J. I. Lunine (Tucson: Univ. Arizona Press), 1253
 Burrows, A., et al. 1997, *ApJ*, 491, 856
 Clampin, M., et al. 2003, *AJ*, 126, 385
 Dent, W. R. F., Walker, H. J., Holland, W. S., & Greaves, J. S. 2000, *MNRAS*, 314, 702
 Draine, B. T., & Lee, H. M. 1984, *ApJ*, 285, 89
 Gomes, R. S. 2003, *Icarus*, 161, 404
 Greaves, J. S., Wyatt, M. C., Holland, W. S., & Dent, W. R. F. 2004, *MNRAS*, 351, L54
 Kalas, P., Liu, M. C., & Matthews, B. C. 2004, *Science*, 303, 1990
 Kenyon, S. J., & Bromley, B. C. 2004, *AJ*, 127, 513
 Kenyon, S. J., Wood, K., Whitney, B. A., & Wolff, M. J. 1999, *ApJ*, 524, L119
 Koen, C., & Eyer, L. 2002, *MNRAS*, 331, 45
 Krist, J., et al. 2004, *ApJ*, in press
 Laor, A., & Draine, B. T. 1993, *ApJ*, 402, 441
 Levison, H. F., & Morbidelli, A. 2003, *Nature*, 426, 419
 Liu, M. 2004, *Science*, 305, 1442
 Takeuchi, T., & Artymowicz, P. 2001, *ApJ*, 557, 990
 Weidenschilling, S. J., & Marzari, F. 1996, *Nature*, 384, 619
 Weinberger, A. J., Becklin, E. E., Schneider, G., Smith, B. A., Lowrance, P. J., Silverstone, M. D., Zuckerman, B., & Terrell, R. J. 1999, *ApJ*, 525, L53
 Weingartner, J. C., & Draine, B. T. 2001, *ApJ*, 548, 296
 Williams, J. P., Najita, J., Liu, M. C., Bottinelli, S., Carpenter, J. M., Hillenbrand, L. A., Meyer, M. R., & Soderblom, D. R. 2004, *ApJ*, 604, 414